

# Pediculosis (lice, head lice, body lice, pubic lice, cooties, crabs)

Last Reviewed: October 2011

## What is pediculosis?

Pediculosis is an infestation of the hairy parts of the body or clothing with the eggs, larvae or adults of lice. The crawling stages of this insect feed on human blood, which can result in severe itching. Head lice are usually located on the scalp, crab lice in the pubic area and body lice along seams of clothing. Body lice travel to the skin to feed and return back to the clothing.

## Who gets pediculosis?

Anyone may become louse infested under suitable conditions of exposure. Pediculosis is easily transmitted from person to person during direct contact. Head lice infestations are frequently found in school settings or institutions. Crab lice infestations can be found among sexually active individuals. Body lice infestation can be found in people living in crowded, unsanitary conditions where clothing is infrequently changed or laundered.

## How is pediculosis spread?

For both head lice and body lice, transmission can occur during direct contact with an infested individual. Sharing of clothing and combs or brushes may also result in transmission of these insects. While other means are possible, crab lice are most often transmitted through sexual contact.

## What are the symptoms of pediculosis?

Usually, the first indication of an infestation is the itching or scratching in the area of the body where the lice feed. Scratching at the back of the head or around the ears should lead to an examination for head louse eggs (nits) on the hair. Itching around the genital area should lead to an examination for crab lice or their eggs. Scratching can be sufficiently intense to result in secondary bacterial infection in these areas.

## How soon do symptoms appear?

It may take as long as two to three weeks or longer for a person to notice the intense itching associated with this infestation.

## For how long is a person able to spread pediculosis?

Pediculosis can be spread as long as lice or eggs remain alive on the infested person or clothing.

## What is the treatment for pediculosis?

Medicated shampoos or cream rinses containing pyrethrins or permethrin are preferred for treating people with head lice. Products containing pyrethrins, permethrin or malathion are available over-the-counter, but those containing lindane or malathion are available only through a physician's prescription. Lindane-based shampoos are not recommended for infants, young children, pregnant women, lactating women, the elderly, persons with HIV or seizure disorders, persons who have very irritated skin or sores where the lindane will be applied and persons who weigh less than 110 pounds. Retreatment after 7 to 10 days is often recommended to ensure that no eggs have survived. Nit combs are available to help remove nits from hair. Dose and duration of shampoo treatment should be followed carefully according to label instructions.

## What can be done to prevent the spread of pediculosis?

Physical contact with infested individuals and their belongings, especially clothing, headgear and bedding, should be avoided. Health education on the life history of lice, proper treatment and the importance of

laundering clothing and bedding in hot water or dry cleaning to destroy lice and eggs is extremely valuable. In addition, regular direct inspection of children for head lice, and when indicated, of body and clothing, particularly of children in schools, institutions, nursing homes and summer camps, is important.

## PEDICULOSIS (Head Lice)

- Pediculosis is defined as an infestation of the hairy parts of the body or clothing with the eggs, larvae, or adults of lice.
- When a student has a confirmed case of pediculosis, parents/guardians will be notified and student must go home with the parent/guardian for treatment.
- Nursing will offer instructions to parents/guardians about effective treatment products, removing nits and cleaning the environment. Treatment to be a pediculocide, such as “Nix”, and combing.
- The student will be allowed to return to school the next day for a reassessment. The student must be brought to the health office by the parent/guardian prior to going to the classroom. The student is not allowed to ride the bus until cleared by the nurse.
- Nursing staff will reassess on a daily basis for live lice and nits for 5 days.
- If live lice or nits are found again, student will return home with parent for treatment with combing only.

## **PEDICULOSIS (Cont.)**

- Re-treatment is required with the pediculocide on the 9<sup>th</sup> day after first treatment. If after the second treatment, student remains with live lice, refer to MD.
- The Clinton County Health Department will be contacted for assistance if necessary.